
Ultimate approximations in nonmonotonic knowledge representation systems

Marc Denecker
Department of Computer Science
K.U.Leeuven
Celestijnenlaan 200A, B-3001 Heverlee
Belgium

Victor W. Marek
Department of Computer Science
University of Kentucky
Lexington, KY, 40506-0046
USA

Mirosław Truszczyński
Department of Computer Science
University of Kentucky
Lexington, KY, 40506-0046
USA

Abstract

We study fixpoints of operators on lattices. To this end we introduce the notion of an approximation of an operator. We order approximations by means of a *precision* ordering. We show that each lattice operator O has a unique most precise or *ultimate* approximation. We demonstrate that fixpoints of this ultimate approximation provide useful insights into fixpoints of the operator O .

We apply our theory to logic programming and introduce the ultimate Kripke-Kleene, well-founded and stable semantics. We show that the ultimate Kripke-Kleene and well-founded semantics are more precise than their standard counterparts. We argue that ultimate semantics for logic programming have attractive epistemological properties and that, while in general they are computationally more complex than the standard semantics, for many classes of theories, their complexity is no worse.

1 INTRODUCTION

Semantics of most knowledge representation languages are defined as collections of interpretations or possible-world structures. The sets of interpretations and possible-world structures, with some natural orderings, form complete lattices. Logic programs, and default and autoepistemic theories determine operators on these lattices. In many cases, semantics of programs and theories are given as fixpoints of these operators. Consequently, an abstract framework of lattices, operators on lattices and their fixpoints has emerged as a powerful tool in investigations of semantics of these logics. Studying semantics of nonmonotonic reasoning systems within an algebraic framework allows us

to eliminate inessential details specific to a particular logic, simplify arguments and find common principles underlying different nonmonotonic formalisms.

The roots of this algebraic approach can be traced back to studies of semantics of logic programs [vEK76, AvE82, Fit85, Prz90] and of applications of lattices and bilattices in knowledge representation [Gin88]. Exploiting the concept of a bilattice and relying on some general properties of operators on lattices and bilattices, Fitting proposed an elegant algebraic treatment of all major 2-, 3- and 4-valued semantics of logic programs [Fit01], that is, the supported-model semantics [Cla78], stable-model semantics [GL88], Kripke-Kleene semantics [Fit85, Kun87] and well-founded semantics [VRS91].

In [DMT00a], we extended Fitting's work to a more abstract setting of the study of fixpoints of lattice operators. Central to our approach is the concept of an approximation of a lattice operator O . An approximation is an operator defined on a certain bilattice (the product of the lattice by itself, with two appropriately defined lattice orderings). Using purely algebraic techniques, for an approximation operator for O we introduced the notion of the stable operator and the concepts of the Kripke-Kleene, well-founded and stable fixpoints, and showed how they provide information about fixpoints of the operator O . In [DMT00a] we noted that our approach generalizes the results described in [Fit01]. We observed that the 4-valued immediate consequence operator \mathcal{T}_P is an approximation operator for the 2-valued immediate consequence operator T_P and showed that all the semantics considered by Fitting can be derived from \mathcal{T}_P by means of the general algebraic constructions that apply to *arbitrary* approximation operators.

In [DMT00b], we applied our algebraic approach to default and autoepistemic logics. Autoepistemic logic was defined by Moore [Moo84] to formalize the knowl-

edge of a rational agent with full introspection capabilities. In Moore’s approach, an autoepistemic theory T defines a characteristic operator D_T on the lattice of all possible-world structures. Fixpoints of D_T (or, to be precise, their theories) are known as *expansions*. In [DMT00b], we proposed for D_T an approximation operator, \mathcal{D}_T , defined on a bilattice of belief pairs (pairs of possible-world structures). Complete fixpoints of \mathcal{D}_T correspond to expansions of T (fixpoints of D_T), the least fixpoint of \mathcal{D}_T provides a constructive approximation to all expansions (by analogy with logic programming, we called it the *Kripke-Kleene fixpoint*). Using general techniques introduced in [DMT00a] we derived from \mathcal{D}_T its *stable* counterpart, the operator \mathcal{D}_T^{st} . Complete fixpoints of \mathcal{D}_T^{st} yield a new semantics of *extensions* for autoepistemic logic. Finally, the least fixpoint of the stable operator results in yet another new semantics, the well-founded semantics for autoepistemic logic (again, called so due to analogies to the well-founded semantics in logic programming), which approximates all extensions.

The same picture emerged in the case of default logic [DMT00b]. For a default theory Δ we defined an operator \mathcal{E}_Δ and characterized all major semantics for default logic in terms of fixpoints of \mathcal{E}_Δ . In particular, the standard semantics of extensions [Rei80] is determined by complete fixpoints of the stable operator \mathcal{E}_Δ^{st} derived from \mathcal{E}_Δ . Our results on autoepistemic and default logics obtained in [DMT00b] allowed us to clarify the issue of their mutual relationship and provided insights into fundamental constructive principles underlying these two modes of nonmonotonic reasoning.

These result prove that the algebraic framework developed in [DMT00a] is an effective tool in studies of semantics of knowledge representation formalisms. It allowed us to establish a comprehensive semantic treatment for nonmonotonic logics and demonstrated that major nonmonotonic systems are closely related. However, the approach, as it was developed, is not entirely satisfactory. It provides no criteria that would allow us to prefer one approximation over another when attempting to define the concept of a stable fixpoint or when approximating fixpoints by means of the Kripke-Kleene or well-founded fixpoints. It does not give us any general indications how to obtain approximations and which approximation to pick. Thus, our theory leaves out a key link in the process of defining and approximating fixpoints of operators on lattices.

In particular, when defining semantics of nonmonotonic formalisms, we *select* an approximation operator, rather than derive it in a principled way. The

approximations used, the bilattice operators \mathcal{T}_P , \mathcal{D}_T and \mathcal{E}_Δ , are not algebraically determined by their corresponding lattice operators T_P , D_T and E_Δ , respectively. Consequently, some programs or theories with the same basic operators have different Kripke-Kleene, well-founded or stable fixpoints associated with them.

We address this problem here. We extend our theory of approximations and introduce the notion of the precision of an approximation. We show that each lattice operator O has a unique *most precise* approximation which we call the *ultimate approximation* of O . Since the ultimate approximation is determined by O , it is well suited for investigations of fixpoints of O . As a result we obtain concepts of ultimate stable fixpoints, the ultimate Kripke-Kleene fixpoint and the ultimate well-founded fixpoint that depend on O only and not on a (possibly arbitrarily) selected approximation to O .

We apply our theory to logic programming, default logic and autoepistemic logic (only the first system is discussed here, due to space limitations). We compare ultimate semantics with the corresponding “standard” semantics of logic programs. In particular, we show that the ultimate Kripke-Kleene and the ultimate well-founded semantics are more precise than the standard Kripke-Kleene and well-founded semantics. This better accuracy comes, however, at a cost. We show that ultimate semantics are in general computationally more complex. On the other hand, we show that for wide classes of theories, including theories likely to occur in practice, the complexity remains the same. Thus, our new semantics may prove useful in computing stable models and default extensions.

The ultimate semantics have also properties that are attractive from the logic perspective. In particular, two programs or theories determining the same basic 2-valued operator have the same ultimate semantics. This property, as we noted, is not true in the standard case.

In summary, our contributions are as follows. We extend the algebraic theory of approximations by providing a principled way of deriving an approximation to a lattice operator. In this way, we obtain concepts of Kripke-Kleene fixpoint, well-founded fixpoint and stable fixpoints that are determined by the operator O and not by the choice of an approximation. In specific contexts of most commonly used nonmonotonic systems we obtain new semantics with desirable logical properties and possible computational applications.

2 PRELIMINARIES

Let $\langle L, \leq \rangle$ be a poset and let A be an operator on L . A poset is *chain-complete* if it contains the least element \perp and if every chain of elements of L has a least upper bound (*lub*) in L . An element x of A is a *pre-fixpoint* of A if $A(x) \leq x$; x is a *fixpoint* of A if $A(x) = x$.

Let A be a monotone operator on a chain-complete poset $\langle L, \leq \rangle$. Let us define a sequence of elements of L by transfinite induction as follows: (1) $c^0 = \perp$; (2) $c^{\alpha+1} = A(c^\alpha)$; (3) $c^\alpha = \text{lub}(\{c^\beta : \beta < \alpha\})$, for a limit ordinal α . One can show that this sequence is well defined, that is has in L its least upper bound and that this least upper bound is the least fixpoint of A ($\text{lfp}(A)$, in symbols). One can also show that the least fixpoint of a monotone operator on a chain-complete poset is the least pre-fixpoint of A . That is, we have $\text{lfp}(A) = \text{glb}(\{x \in L : A(x) \leq x\})$. Monotone operators on chain-complete posets and their fixpoints and pre-fixpoints are discussed in [Mar76].

A lattice is a poset $\langle L, \leq \rangle$ such that $L \neq \emptyset$ and every pair of elements $x, y \in L$ has a unique greatest lower bound and least upper bound. A lattice is *complete* if its *every* subset has a greatest lower bound and a least upper bound. In particular, a complete lattice has a least and a greatest element denoted by \perp and \top , respectively.

For any two elements $x, y \in L$, we define $[x, y] = \{z \in L : x \leq z \leq y\}$. If $\langle L, \leq \rangle$ is a complete lattice and $x \leq y$, then $\langle [x, y], \leq \rangle$ is a complete lattice, too.

Let $\langle L, \leq \rangle$ be a complete lattice. By the *product bilattice* [Gin88] of $\langle L, \leq \rangle$ we mean the set $L^2 = L \times L$ with the following two orderings \leq_p and \leq :

1. $(x, y) \leq_p (x', y')$ if $x \leq x'$ and $y' \leq y$
2. $(x, y) \leq (x', y')$ if $x \leq x'$ and $y \leq y'$.

Both orderings are complete lattice orderings for L^2 . However, in this paper we are mostly concerned with the ordering \leq_p .

An element $(x, y) \in L^2$ is *consistent* if $x \leq y$. We can think of a consistent element $(x, y) \in L^2$ as an *approximation* to every $z \in L$ such that $x \leq z \leq y$. With this interpretation in mind, the ordering \leq_p , when restricted to consistent elements, can be viewed as a *precision* ordering. Consistent pairs that are “higher” in the ordering \leq_p provide tighter approximations. Maximal consistent elements with respect to \leq_p are pairs of the form (x, x) . We call approximations of the form (x, x) — *exact*.

We denote the set of all consistent pairs in L^2 by L^c . The set $\langle L^c, \leq_p \rangle$ is not a lattice. It is, however, chain-

complete. Indeed, the element (\perp, \top) is the least element in L^c and the following result shows that every chain in L^c has (in L^c) the least upper bound.

Proposition 2.1 *Let L be a complete lattice. If $\{(a^\alpha, b^\alpha)\}_\alpha$ is a chain of elements in $\langle L^c, \leq_p \rangle$ then $\text{lub}(\{a^\alpha\}_\alpha) \leq \text{glb}(\{b^\alpha\}_\alpha)$ and $(\text{lub}(\{a^\alpha\}_\alpha), \text{glb}(\{b^\alpha\}_\alpha)) = \text{lub}_{\leq_p}(\{(a^\alpha, b^\alpha)\}_\alpha)$.*

It follows that every \leq_p -monotone operator on L^c has a least fixpoint.

3 PARTIAL APPROXIMATIONS

For an operator $A : L^c \rightarrow L^c$, we denote by A^1 and A^2 its projections to the first and second coordinates, respectively. Thus, for every $(x, y) \in L^c$, we have $A(x, y) = (A^1(x, y), A^2(x, y))$. An operator $A : L^c \rightarrow L^c$ is a *partial approximation* operator if it is \leq_p -monotone and if for every $x \in L$, $A^1(x, x) = A^2(x, x)$. We denote the set of all partial approximation operators on L^c by $\text{Appx}(L^c)$. Let $A \in \text{Appx}(L^c)$. Since A is \leq_p -monotone and L^c is chain-complete, A has a least fixpoint, called the *Kripke-Kleene fixpoint* of A ($\text{KK}(A)$, in symbols). Directly from the definition, it follows that $\text{KK}(A)$ approximates all fixpoints of A .

If $A \in \text{Appx}(L^c)$ and $O : L \rightarrow L$ is an operator on L such that $A(x, x) = (O(x), O(x))$ then we say that A is a *partial approximation of O* . We denote the set of all partial approximations of O by $\text{Appx}(O)$. If A is a partial approximation of O then $x \in L$ is a fixpoint of O if and only if (x, x) is a fixpoint of A . Thus, for every fixpoint x of O , we have $\text{KK}(A) \leq_p (x, x)$ or, equivalently, $\text{KK}^1(A) \leq x \leq \text{KK}^2(A)$, where $\text{KK}^1(A)$ and $\text{KK}^2(A)$ are the two components of the pair $\text{KK}(A)$.

Operators from $\text{Appx}(L^c)$ describe ways to revise consistent approximations. Of particular interest are those situations when the revision of an approximation leads to another one that is at least as accurate. Let A be an operator on L^c . We call an approximation (a, b) *A-reliable* if $(a, b) \leq_p A(a, b)$.

Proposition 3.1 *Let L be a complete lattice and $A \in \text{Appx}(L^c)$. If $(a, b) \in L^c$ is A-reliable then, for every $x \in [\perp, b]$, $A^1(x, b) \in [\perp, b]$ and, for every $x \in [a, \top]$, $A^2(a, x) \in [a, \top]$.*

Proof: Let $x \in [\perp, b]$. Then $(x, b) \leq_p (b, b)$. By the \leq_p -monotonicity of A ,

$$A^1(x, b) \leq A^1(b, b) = A^2(b, b) \leq A^2(a, b) \leq b.$$

The last inequality follows from the fact that (a, b) is *A-reliable*. The second part of the assertion can be proved in a similar manner. \square

This proposition implies that for every A -reliable pair (a, b) , the restrictions of $A^1(\cdot, b)$ to $[\perp, b]$ and $A^2(a, \cdot)$ to $[a, \top]$ are in fact operators on $[\perp, b]$ and $[a, \top]$, respectively. Moreover, they are \leq -monotone operators on the posets $([\perp, b], \leq)$ and $([a, \top], \leq)$. Since $([\perp, b], \leq)$ and $([a, \top], \leq)$ are complete lattices, the operators $A^1(\cdot, b)$ and $A^2(a, \cdot)$ have least fixpoints in the lattices $([\perp, b], \leq)$ and $([a, \top], \leq)$, respectively. We define:

$$b^{A\downarrow} = \text{lfp}(A^1(\cdot, b)) \quad \text{and} \quad a^{A\uparrow} = \text{lfp}(A^2(a, \cdot)).$$

We call the mapping $(a, b) \mapsto (b^{A\downarrow}, a^{A\uparrow})$, defined on the set of A -reliable elements of L^c , the *stable revision operator for A* . When A is clear from the context, we will drop the reference to A from the notation.

Directly from the definition of the stable revision operator it follows that for every A -reliable pair, $b^\downarrow \leq b$ and $a \leq a^\uparrow$.

The stable revision operator for $A \in \text{Appx}(L^c)$ is crucial. It allows us to distinguish an important subclass of the class of all fixpoints of A . Let L be a complete lattice and let $A \in \text{Appx}(L^c)$. We say that $(x, y) \in L^c$ is a *stable fixpoint* of A if (x, y) is A -reliable and is a fixpoint of the stable revision operator (that is, $x = x^\downarrow$ and $y = y^\uparrow$). By the A -reliability of (x, y) , the second requirement is well defined.

Stable fixpoints of an operator are, in particular, its fixpoints.

Proposition 3.2 *Let L be a complete lattice and let $A \in \text{Appx}(L^c)$. If (x, y) is a stable fixpoint of A then (x, y) is a fixpoint of A .*

Proof: Since (x, y) is stable, $x = \text{lfp}(A^1(\cdot, y))$. In particular, $x = A^1(x, y)$. Similarly, $y = A^2(x, y)$. \square

Let O be an operator on a complete lattice L and let $A \in \text{Appx}(O)$. We say that x is an *A -stable fixpoint* of O if (x, x) is a stable fixpoint of A . The notation is justified. Indeed, it follows from Proposition 3.2 and our earlier remarks that every stable fixpoint of O is, in particular, a fixpoint of O .

The notion of A -reliability is not strong enough to guarantee desirable properties of the stable revision operator. In particular, if $(a, b) \in L^c$ is A -reliable, it is not true in general that $(b^\downarrow, a^\uparrow)$ is consistent nor that $(a, b) \leq_p (b^\downarrow, a^\uparrow)$. There is, however, a class of A -reliable pairs for which both properties hold. An A -reliable approximation (a, b) is *A -prudent* if $a \leq b^\downarrow$. We note that every stable fixpoint of A is A -prudent. We will now prove several basic properties of A -prudent approximations.

Proposition 3.3 *Let L be a complete lattice, $A \in \text{Appx}(L^c)$ and $(a, b) \in L^c$ be A -prudent. Then, $(b^\downarrow, a^\uparrow)$ is consistent, A -reliable and A -prudent and $(a, b) \leq_p (b^\downarrow, a^\uparrow)$.*

Proof: By the definition of b^\downarrow and a^\uparrow we have that $b^\downarrow \leq b$ and $a \leq a^\uparrow$. Moreover, since (a, b) is A -prudent, it follows that $a \leq b^\downarrow$.

Next, since (a, b) is A -reliable, it follows that $a \leq b$ and $A^2(a, b) \leq b$. Thus, b is a pre-fixpoint of $A^2(a, \cdot)$. Consequently, $a^\uparrow \leq b$ (as a^\uparrow is the least fixpoint of $A^2(a, \cdot)$). Hence, $(a, b) \leq_p (b^\downarrow, a^\uparrow)$.

By the \leq_p -monotonicity of A we obtain:

$$A^1(a^\uparrow, b) \leq A^1(a^\uparrow, a^\uparrow) = A^2(a^\uparrow, a^\uparrow) \leq A^2(a, a^\uparrow) = a^\uparrow.$$

It follows that a^\uparrow is a pre-fixpoint of the operator $A^1(\cdot, b)$. Thus, $b^\downarrow = \text{lfp}(A^1(\cdot, b)) \leq a^\uparrow$ and so, $(b^\downarrow, a^\uparrow)$ is consistent.

Let us now observe that $b^\downarrow = A^1(b^\downarrow, b) \leq A^1(b^\downarrow, a^\uparrow)$. Similarly, $a^\uparrow = A^2(a, a^\uparrow) \geq A^2(b^\downarrow, a^\uparrow)$. Thus, the pair $(b^\downarrow, a^\uparrow)$ is reliable.

Lastly, we note that for every $x \in [\perp, a^\uparrow]$, $A^1(x, b) \leq A^1(x, a^\uparrow) \leq a^\uparrow$ (the last inequality follows by the A -reliability of $(b^\downarrow, a^\uparrow)$). Hence, $b^\downarrow = \text{lfp}(A^1(\cdot, b)) \leq \text{lfp}(A^1(\cdot, a^\uparrow))$ and, consequently, $(b^\downarrow, a^\uparrow)$ is A -prudent. \square

Let us observe that an A -reliable pair (a, b) is revised by an operator A into a more accurate approximation $A(a, b)$. An A -prudent pair (a, b) can be revised “even more”. Namely, it is easy to see that $A^1(a, b) \leq A^1(b^\downarrow, b) = b^\downarrow$ and $a^\uparrow = A^2(a, a^\uparrow) \leq A^2(a, b)$. Thus, $A(a, b) \leq_p (b^\downarrow, a^\uparrow)$. In other words, $(b^\downarrow, a^\uparrow)$ is indeed at least as precise revision of (a, b) as $A(a, b)$ is.

The stable revision operator satisfies a certain monotonicity property.

Proposition 3.4 *Let L be a complete lattice and let $A \in \text{Appx}(L^c)$. If $(a, b) \in L^c$ is A -reliable, $(c, d) \in L^c$ is A -prudent and if $(a, b) \leq_p (c, d)$, then $(b^\downarrow, a^\uparrow) \leq_p (d^\downarrow, c^\uparrow)$.*

Proof: Clearly, we have $d^\downarrow \leq c^\uparrow \leq d \leq b$. By the \leq_p -monotonicity of A , it follows that $A^1(d^\downarrow, b) \leq A^1(d^\downarrow, d) = d^\downarrow$. Thus, d^\downarrow is a pre-fixpoint of $A^1(\cdot, b)$. Since b^\downarrow is the least fixpoint of $\text{lfp}(A^1(\cdot, b))$, it follows that $b^\downarrow \leq d^\downarrow$.

It remains to prove that $c^\uparrow \leq a^\uparrow$. Let $u = \text{glb}(a^\uparrow, d^\downarrow)$. By Proposition 3.3, $(c, d) \leq_p (d^\downarrow, c^\uparrow)$. Since $(a, b) \leq_p (c, d)$, it follows that $a \leq d^\downarrow$. Further, by the A -reliability of (a, b) and (c, d) , we have $a \leq a^\uparrow$ and

$d^\downarrow \leq d$. Thus, $a \leq u \leq a^\uparrow$ and $u \leq d^\downarrow \leq d$. Consequently,

$$A^1(u, d) \leq A^1(u, u) = A^2(u, u) \leq A^2(a, a^\uparrow) = a^\uparrow$$

and

$$A^1(u, d) \leq A^1(d^\downarrow, d) = d^\downarrow.$$

It follows that $A^1(u, d) \leq glb(a^\uparrow, d^\downarrow) = u$. In particular, u is a pre-fixpoint of $A^1(\cdot, d)$. Since d^\downarrow is the least fixpoint of $A^1(\cdot, d)$, $d^\downarrow \leq u$. Hence, $d^\downarrow \leq a^\uparrow$.

We now have $a \leq c \leq d^\downarrow \leq a^\uparrow$ (the first inequality follows from the assumption $(a, b) \leq (c, d)$, the second one follows by Proposition 3.3 from the assumption that (c, d) is A -prudent). Thus, $a \leq c \leq a^\uparrow$ and the \leq_p -monotonicity of A implies

$$A^2(c, a^\uparrow) \leq A^2(a, a^\uparrow) = a^\uparrow.$$

Hence, a^\uparrow is a pre-fixpoint of $A^2(c, \cdot)$. Since c^\uparrow is the least fixpoint of $A^2(c, \cdot)$, it follows that $c^\uparrow \leq a^\uparrow$. \square

Since stable fixpoints are prudent, we obtain the following corollary.

Corollary 3.5 *Let L be a complete lattice, $A \in \text{Appx}(L^c)$ and let $(c, d) \in L^c$ be a stable fixpoint of A . If $(a, b) \in L^c$ is A -reliable and $(a, b) \leq_p (c, d)$ then $(b^\downarrow, a^\uparrow) \leq_p (c, d)$. \square*

The next result states that the limit of a chain of A -prudent pairs is A -prudent.

Proposition 3.6 *Let L be a complete lattice, $A \in \text{Appx}(L^c)$ and let $\{(a^\alpha, b^\alpha)\}_\alpha$ be a chain of A -prudent pairs from L^c . Then, $\text{lub}(\{(a^\alpha, b^\alpha)\}_\alpha)$ is A -prudent.*

Proof: Let us set $a^\infty = \text{lub}(\{a^\alpha\}_\alpha)$ and $b^\infty = glb(\{b^\alpha\}_\alpha)$. By Proposition 2.1, (a^∞, b^∞) is consistent and $(a^\infty, b^\infty) = \text{lub}(\{(a^\alpha, b^\alpha)\}_\alpha)$. Let us now observe that, by A -reliability of (a^α, b^α) and \leq_p -monotonicity of A , we have $(a^\alpha, b^\alpha) \leq_p A(a^\alpha, b^\alpha) \leq_p A(a^\infty, b^\infty)$. Thus, $(a^\infty, b^\infty) = \text{lub}(\{(a^\alpha, b^\alpha)\}_\alpha) \leq A(a^\infty, b^\infty)$. It follows that (a^∞, b^∞) is A -reliable.

The A -reliability of (a^∞, b^∞) implies, in particular, that for every $x \in [\perp, b^\infty]$, $A^1(x, b^\infty) \leq b^\infty$. Thus, by \leq_p -monotonicity of A , for every $x \in [\perp, b^\infty]$

$$A^1(x, b^\alpha) \leq A^1(x, b^\infty) \leq b^\infty.$$

Hence, pre-fixpoints of $A^1(\cdot, b^\infty)$ are prefixpoints of $A^1(\cdot, b^\alpha)$ and, consequently,

$$\text{lfp}(A^1(\cdot, b^\alpha)) \leq \text{lfp}(A^1(\cdot, b^\infty)).$$

Since (a^α, b^α) is A -prudent, we have that $a^\alpha \leq \text{lfp}(A^1(\cdot, b^\alpha))$. Thus, for arbitrary α , $a^\alpha \leq \text{lfp}(A^1(\cdot, b^\infty))$ and, consequently, $a^\infty \leq \text{lfp}(A^1(\cdot, b^\infty))$. It follows that (a^∞, b^∞) is A -prudent. \square

We will now prove that the set of all stable fixpoints of an operator has a least element (in particular, it is not empty). To this end, we define a sequence $\{(a^\alpha, b^\alpha)\}_\alpha$ of elements of L^c by transfinite induction:

1. $(a^0, b^0) = (\perp, \top)$
2. If $\alpha = \beta + 1$, we define $a^\alpha = b^{\beta\downarrow}$ and $b^\alpha = a^{\beta\uparrow}$
3. If α is a limit ordinal, we define $(a^\alpha, b^\alpha) = \text{lub}(\{(a^\beta, b^\beta) : \beta < \alpha\})$.

Theorem 3.7 *The sequence $\{(a^\alpha, b^\alpha)\}_\alpha$ is well defined, \leq_p -monotone and its limit is the least stable fixpoint of a partial approximation operator A .*

Proof: It is obvious that (\perp, \top) is A -prudent. Thus, by the transfinite induction it follows that each element in the sequence is well defined and A -prudent (Propositions 3.3 and 3.6 settle the cases of successor ordinals and limit ordinals, respectively). In the same way, one can establish the \leq_p -monotonicity of the sequence.

Let $(a^\infty, b^\infty) = \text{lub}(\{(a^\beta, b^\beta)\}_\alpha)$. By Proposition 3.6, (a^∞, b^∞) is A -prudent. Thus, (a^∞, b^∞) is A -reliable. Moreover, we have $a^\infty = (b^\infty)^\downarrow$ and $b^\infty = (a^\infty)^\uparrow$. Thus, (a^∞, b^∞) is a stable fixpoint of A . Further, it is easy to see by transfinite induction and Corollary 3.5 that (a^∞, b^∞) approximates all stable fixpoints of A . Thus, it is the least stable fixpoint of A . \square

We call this least stable fixpoint the *well-founded fixpoint* of A and denote it by $\text{WF}(A)$. The well-founded fixpoint approximates all stable fixpoints of A . In particular, it approximates all A -stable fixpoints of the operator O . That is, for every A -stable fixpoint x of O , $\text{WF}(A) \leq_p (x, x)$ or, equivalently, $\text{WF}^1(A) \leq x \leq \text{WF}^2(A)$, where $\text{WF}^1(A)$ and $\text{WF}^2(A)$ are the two components of the pair $\text{WF}(A)$. Moreover, the well-founded fixpoint is more precise than the Kripke-Kleene fixpoint: for $A \in \text{Appx}(O)$, $\text{KK}(A) \leq_p \text{WF}(A)$.

In [DMT00b, DMT00a], we showed that when applied to appropriately chosen approximation operators in logic programming, default logic and autoepistemic logic, these algebraic concepts of fixpoints, stable fixpoints, the Kripke-Kleene fixpoint and the well-founded fixpoint provide all major semantics for these nonmonotonic systems and allow us to understand their interrelations.

We need to emphasize that the concept of a partial approximation introduced here is different from the concept of *approximation* introduced in [DMT00a]. The

latter notion is defined as an operator of the whole bilattice L^2 . That choice was motivated by our search for generality and potential applications of inconsistent fixpoints in situations when we admit a possibility of some statements being overdefined. While different, both approaches are very closely related¹.

4 ULTIMATE APPROXIMATIONS

Partial approximations in $Appx(L^c)$ can be ordered. Let $A, B \in Appx(L^c)$. We say that A is *less precise* than B ($A \leq_p B$, in symbols) if for each pair $(x, y) \in L^c$, $A(x, y) \leq_p B(x, y)$. It is easy to see that if $A \leq_p B$ then there is an operator O on the lattice L such that $A, B \in Appx(O)$.

Lemma 4.1 *Let L be a complete lattice and $A, B \in Appx(L^c)$. If $A \leq_p B$ and $(a, b) \in L^c$ is A -prudent then (a, b) is B -prudent and $(b^{A\downarrow}, a^{A\uparrow}) \leq_p (b^{B\downarrow}, a^{B\uparrow})$.*

Proof: Clearly, $(a, b) \leq_p A(a, b) \leq B(a, b)$. Thus, (a, b) is B -reliable.

For each pre-fixpoint $x \leq b$ of $B^1(\cdot, b)$, $A^1(x, b) \leq B^1(x, b) \leq x$. Consequently, x is a prefixpoint of $A^1(\cdot, b)$. It follows that $b^{A\downarrow} \leq b^{B\downarrow}$. Since $a \leq b^{A\downarrow}$, $a \leq b^{B\downarrow}$. Thus (a, b) is B -prudent.

Likewise, we can prove that any pre-fixpoint of $A^2(a, \cdot)$ is a prefixpoint of $B^2(a, \cdot)$, and consequently, $a^{B\uparrow} \leq a^{A\uparrow}$. Since also $b^{A\downarrow} \leq b^{B\downarrow}$, it follows that $(b^{A\downarrow}, a^{A\uparrow}) \leq_p (b^{B\downarrow}, a^{B\uparrow})$. \square

More precise approximation have more precise Kripke-Kleene and well-founded fixpoints.

Theorem 4.2 *Let O be an operator on a complete lattice L . Let $A, B \in Appx(O)$. If $A \leq_p B$ then $KK(A) \leq_p KK(B)$ and $WF(A) \leq_p WF(B)$.*

Proof: Let us denote by $\{(a_A^\alpha, b_A^\alpha)\}_\alpha$ the sequence of elements of $\langle L^c, \leq_p \rangle$ obtained by iterating the operator A over (\perp, \top) . The sequence $\{(a_B^\alpha, b_B^\alpha)\}_\alpha$ is defined in the same way. Since $A \leq_p B$, it follows by an easy induction that for every ordinal α , $(a_A^\alpha, b_A^\alpha) \leq_p (a_B^\alpha, b_B^\alpha)$. Since $KK(A)$ is the limit of the sequence $\{(a_A^\alpha, b_A^\alpha)\}_\alpha$ and $KK(B)$ is the limit of the sequence $\{(a_B^\alpha, b_B^\alpha)\}_\alpha$, it follows that $KK(A) \leq_p KK(B)$.

To prove the second part of the assertion, we will now assume that the sequences $\{(a_A^\alpha, b_A^\alpha)\}_\alpha$ and $\{(a_B^\alpha, b_B^\alpha)\}_\alpha$ denote the sequences used in the definition of the well-founded fixpoints of A and B , respec-

tively. To prove the assertion we will now show that for every ordinal α , $(a_A^\alpha, b_A^\alpha) \leq_p (a_B^\alpha, b_B^\alpha)$.

Clearly, $(a_A^0, b_A^0) \leq_p (a_B^0, b_B^0)$. Let us assume that $\alpha = \beta + 1$ and that $(a_A^\beta, b_A^\beta) \leq_p (a_B^\beta, b_B^\beta)$. Since (a_A^β, b_A^β) is A -prudent, Lemma 4.1 entails that it is B -prudent and

$$(a_A^\alpha, b_A^\alpha) = ((b_A^\beta)^{A\downarrow}, (a_A^\beta)^{A\uparrow}) \leq_p ((b_A^\beta)^{B\downarrow}, (a_A^\beta)^{B\uparrow}).$$

By Proposition 3.4,

$$((b_A^\beta)^{B\downarrow}, (a_A^\beta)^{B\uparrow}) \leq_p ((b_B^\beta)^{B\downarrow}, (a_B^\beta)^{B\uparrow}) = (a_B^\alpha, b_B^\alpha).$$

The case of the limit ordinal α is straightforward.

Since $WF(A)$ and $WF(B)$ are the limits of the sequences $\{(a_A^\alpha, b_A^\alpha)\}_\alpha$ and $\{(a_B^\alpha, b_B^\alpha)\}_\alpha$, respectively, the second part of the assertion follows. \square

The next result shows that as the precision of an approximation grows, all exact fixpoints and exact stable fixpoints are preserved.

Theorem 4.3 *Let O be an operator on a complete lattice L . Let $A, B \in Appx(O)$. If $A \leq_p B$ then every exact fixpoint of A is an exact fixpoint of B , and every exact stable fixpoint of A (that is, an A -stable fixpoint of O) is also an exact stable fixpoint of B (that is, a B -stable fixpoint of O).*

Proof: Since for every $x \in L$, $A(x, x) = B(x, x) = (O(x), O(x))$, the first part of the assertion follows. Let us now assume that (x, x) is an exact stable fixpoint of A . In particular, it follows that (x, x) is a fixpoint of A and is A -prudent. By Lemma 4.1, (x, x) is B -prudent and $(x, x) \leq_p (x^{B\downarrow}, x^{B\uparrow})$. The latter pair is consistent (Proposition 3.3). Consequently, (x, x) is $(x^{B\downarrow}, x^{B\uparrow})$ and hence x is an exact stable fixpoint of B . \square

Non-exact fixpoints are not preserved, in general. Let us consider two partial approximations A and B such that $A \leq_p B$. Let us also assume that $WF(A) <_p WF(B)$ (that is, A has a strictly less precise well-founded fixpoint than B). Then, clearly, $WF(A)$ is no longer a stable fixpoint of B . Thus, fixpoints of A may disappear when we move on to a more precise approximation B .

More precise approximations of a non-monotone operator O yield more precise well-founded fixpoints and additional exact stable fixpoints. The natural question is whether there exists an *ultimate approximation* of O , that is, a partial approximation most precise with respect to the ordering \leq_p . Such approximation would have a most precise Kripke-Kleene and well-founded

¹We will include a detailed discussion of the relationship between the two approaches in the full version of the paper.

fixpoint and a largest set of exact stable fixpoints. We will show that the answer to this key question is positive. Such ultimate approximation, being a distinguished object in the collection of all approximations can be viewed as determined by O . Consequently, fixpoints of the ultimate approximation of O (including stable, Kripke-Kleene and well-founded fixpoints) can be regarded as determined by O and can be associated with it.

We start by providing a non-constructive argument for the existence of ultimate approximations. Let us note that the set $Appx(O)$ is not empty. Indeed, let us define $A_O(x, y) = (O(x), O(x))$, if $x = y$, and $A_O(x, y) = (\perp, \top)$, otherwise. It is easy to see that $A_O \in Appx(O)$ and that it is the least precise element in $Appx(O)$. Next, we observe that $Appx(O)$ with the ordering \leq_p is a complete lattice, as the set $Appx(O)$ is closed under the operations of taking greatest lower bounds and least upper bounds. It follows that $Appx(O)$ has a greatest element (most precise approximation). We call this partial approximation the *ultimate approximation* of O and denote it by U_O .

We call the Kripke-Kleene and the well-founded fixpoints of U_O , the *ultimate Kripke-Kleene* and the *ultimate well-founded fixpoint* of O . We denote them by $KK(O)$ and $WF(O)$, respectively. We call a stable fixpoint of U_O an *ultimate partial stable fixpoint* of O . We refer to an *exact stable fixpoint* of U_O as an *ultimate stable fixpoint* of O . Exact fixpoints of all partial approximations are the same and correspond to fixpoints of O . Thus, there is no need to introduce the concept of an ultimate exact fixpoint of O . We have the following corollary to Theorems 4.2 and 4.3.

Corollary 4.4 *Let O be an operator on a complete lattice L . For every $A \in Appx(O)$, $KK(A) \leq_p KK(U_O)$, $WF(A) \leq_p WF(U_O)$ and every A -stable fixpoint of O is an ultimate stable fixpoint of O .*

We will now provide a constructive characterization of the notion. To state the result, for every $x, y \in L$ such that $x \leq y$, we define $O([x, y]) = \{O(z) : z \in [x, y]\}$.

Theorem 4.5 *Let O be an operator on a complete lattice L . Then, for every $(x, y) \in L^c$, $U_O(x, y) = (glb(O([x, y])), lub(O([x, y])))$.*

Proof: We define an operator $C : L^c \rightarrow L^2$ by setting

$$C(x, y) = (glb(O([x, y])), lub(O([x, y]))).$$

First, let us notice that since $glb(O([x, y])) \leq lub(O([x, y]))$, the operator C maps L^c into L^c . Moreover, it is easy to see that C is \leq_p -monotone. Lastly,

since $O([x, x]) = \{O(x)\}$,

$$glb(O([x, x])) = lub(O([x, x])) = O(x).$$

and, consequently, $C(x, x) = (O(x), O(x))$. Thus, it follows that C is a partial approximation of O . Since U_O is the most precise approximation, we have $C \leq_p U_O$.

On the other hand, $U_O(x, y) \leq (O(z), O(z))$ for every $z \in [x, y]$. Therefore $U_O^1(x, y) \leq_p O(z)$ for all $z \in [x, y]$ and thus $U_O^1(x, y) \leq glb(O([x, y]))$. Similarly, $lub(O([x, y])) \leq U_O^2(x, y)$. Since $x \leq y$ are arbitrary, $U_O \leq_p C$, as desired. \square

With this result we obtain an explicit characterization of ultimate stable fixpoints of an operator O .

Corollary 4.6 *Let L be a complete lattice. An element $x \in L$ is an ultimate stable fixpoint of an operator $O : L \rightarrow L$ if and only if x is the least fixpoint of the operator $glb(O([\cdot, x]))$ regarded as an operator on $[\perp, x]$.*

We conclude this section by describing ultimate approximations for monotone and antimonotone operators on L .

Proposition 4.7 *If O is a monotone operator on a complete lattice L then for every $(x, y) \in L^c$, $U_O(x, y) = (O(x), O(y))$. If O is antimonotone then for every $(x, y) \in L^c$, $U_O(x, y) = (O(y), O(x))$.*

Proof: By Theorem 4.5,

$$U_O(x, y) = (glb(O([x, y])), lub(O([x, y]))).$$

Now, it is easy to see that if O is monotone, then $glb(O([x, y])) = O(x)$ and $lub(O([x, y])) = O(y)$. If O is antimonotone, then $glb(O([x, y])) = O(y)$ and $lub(O([x, y])) = O(x)$. The proposition follows. \square

Using the results from [DMT00a] and Proposition 4.7 we now obtain the following corollary.

Corollary 4.8 *Let O be an operator on a complete lattice L . If O is monotone, then the least fixpoint of O is the ultimate well-founded fixpoint of O and the unique ultimate stable fixpoint of O . If O is antimonotone, then $KK(O) = WF(O)$ and every fixpoint of O is an ultimate stable fixpoint of O .*

5 ULTIMATE SEMANTICS FOR LOGIC PROGRAMMING

The basic operator in logic programming is the one-step provability operator T_P introduced in [vEK76]. It

is defined on the lattice of all interpretations. This lattice consists of subsets of the set of all atoms appearing in P and is ordered by inclusion (we identify truth assignments with subsets of atoms that are assigned the value \mathbf{t}).

Let P be a logic program. We denote by U_P the ultimate approximation operator for the operator T_P . By specializing Theorem 4.5 to the operator T_P we obtain that for every two interpretations $I \subseteq J$,

$$U_P(I, J) = (glb(T_P([I, J])), lub(T_P([I, J]))).$$

Replacing the ultimate approximation operator U_O in the definitions of ultimate Kripke-Kleene, well-founded and stable fixpoints with U_P results in the corresponding notions of ultimate Kripke-Kleene, well-founded and stable models (semantics) of a program P .

We are now in a position to discuss commonsense reasoning intuitions underlying abstract algebraic concepts of ultimate approximation and its fixpoints. Let us consider two interpretations I and J such that $I \subseteq J$. We interpret I as a current lower bound and J as a current upper bound on the set of atoms that are true (under P). Thus, I specifies atoms that are definitely true, while J specifies atoms that are possibly true. Arguably, if an atom p is derived by applying the operator T_P to every interpretation $K \in [I, J]$, it can safely be assumed to be true (in the context of the knowledge represented by I and J). Thus, the set $I' = glb(T_P([I, J]))$ can be viewed as a revision of I .

Similarly, since every interpretation $K \in [I, J]$ must be regarded as possible according to the pair (I, J) of conservative and liberal estimates, an atom might possibly be true if it can be derived by the operator T_P from at least one interpretation in $[I, J]$. Thus, the set $J' = lub(T_P([I, J]))$, consisting of all such atoms, can be regarded as a revision of J . Clearly, $(I', J') = U_P(I, J)$ and, consequently, U_P can be viewed as a way to revise our knowledge about the logical values of atoms as determined by a program P from (I, J) to (I', J') .

By iterating U_P starting at (\perp, \top) , we obtain the ultimate Kripke-Kleene model of P as an approximation that cannot be further improved by applying U_P . The ultimate Kripke Kleene model of P approximates all fixpoints of U_P and, in particular, all supported models of P . Often, however, the Kripke-Kleene model is too weak as we are commonly interested in those (partial) models of P that satisfy some minimality or groundedness conditions. These requirements are satisfied by ultimate stable models and, in particular, by the ultimate well-founded model of P .

When constructing the ultimate well-founded model, we start by assuming no knowledge about the status of atoms: no atom is known true and all atoms are assumed possible. Our goal is to improve on these bounds.

To improve on the lower bound, we proceed as follows. Our current knowledge does not preclude any interpretation and all of them (the whole segment $[\perp, \top]$) need to be taken into account. If some atom p can be derived by applying the operator T_P to each element of $[\perp, \top]$ then, arguably, p could be accepted as definitely true. The set of all these atoms is exactly $glb(T_P([\perp, \top]))$. So, this set, say I_1 , can be taken as a safe new lower bound, giving a smaller interval $[I_1, \top]$ of possible interpretations. We now repeat the same process and obtain a new lower bound, say I_2 , consisting of those atoms that can be derived from every interpretation in $[I_1, \top]$. It is given by $I_2 = glb(T_P([I_1, \top]))$. Clearly, I_2 improves on I_1 . We iterate this process until a fixpoint is reached. This fixpoint, say I^1 , consists of all these atoms for which there is a *constructive* argument that they are true, given that no atoms are known to be false (all atoms are possible). Thus, it provides a safe lower bound for the set of atoms the program should specify as true.

The reasoning for revising the upper bound is different. The goal is to make false all atoms for which there cannot be a constructive argument that they are true. Let us consider an interpretation J such that for every $K \in [\perp, J]$, $T_P(K) \in [\perp, J]$, or equivalently, $lub(T_P([\perp, J])) \subseteq J$. An atom $p \notin J$ (false in J) cannot be made true by applying T_P to any element in the segment $[\perp, J]$. In order to derive p by means of T_P , some atoms that are false in J would have to be made true. That, however, would mean that p is not grounded and could be assumed to be false. Thus, each such interpretation J represents an upper estimate on what is possible (its complement gives a lower estimate on what is false) under the assumption that no atom is known to be true yet. It turns out that there is a least interpretation, say J^1 such that $lub(T_P([\perp, J^1])) \subseteq J^1$ and it can be constructed in a bottom up way by iterating the operator $lub(T_P([\perp, \cdot]))$. This interpretation can be taken as a safe lower bound on what is false (given that no atom is known to be true).

The pair (I^1, J^1) is the first improvement on (\perp, \top) . It is precisely the pair produced by the first iteration of the general well-founded fixpoint definition given earlier. It can now be used, in place of (\perp, \top) , to obtain an even more refined estimate, (I^2, J^2) and the process continues until the fixpoint is reached. The resulting pair is the ultimate well-founded model of P .

This discussion demonstrates that abstract algebraic concepts of ultimate approximations can be given a sound intuitive account.

We will now discuss the properties of the ultimate semantics for logic programs.

Theorem 5.1 *Let P, P' be two programs such that $T_P = T_{P'}$. Then, the ultimate well-founded models and ultimate stable models of P and P' coincide.*

Proof: Theorem 4.5 implies that $U_P = U_{P'}$. But then all fixpoints of U_P and $U_{P'}$ coincide. Thus, the result follows. \square

This assertion does not hold for the (standard) well-founded and stable models. For instance, let $P_1 = \{p \leftarrow p, p \leftarrow \neg p\}$ and $P_2 = \{p \leftarrow \cdot\}$. Clearly, $T_{P_1} = T_{P_2}$. However, P_2 has a stable model, $\{p\}$, while P_1 has no stable models. Furthermore, p is true in the well-founded model of P_2 and unknown in the well-founded model of P_1 .

Another appealing property is that the ultimate well-founded model of a program P with monotone operator T_P is the least fixpoint of this operator (the least model of P). This is a corollary of Proposition 4.8. It is not satisfied by the standard well-founded semantics, as shown by the program P_1 .

In many cases, the ultimate well-founded semantics coincides with the standard well-founded semantics. A consequence of Corollary 4.4 is that if the well-founded model of a program is two-valued, then it coincides with the ultimate well-founded model. Thus, we have the following result dealing with the classes of Horn and weakly stratified programs [Prz90]:

Proposition 5.2 *If a logic program P is a Horn program or a (weakly) stratified program, then its ultimate well-founded semantics coincides with the standard well-founded semantics.*

Proof: Let P be a Horn program or a weakly stratified program (the argument is the same). Let WF_P be the well-founded model of P . Let T_P be the van Emden-Kowalski operator for P , and let \mathcal{T}_P be the corresponding 3-valued operator [Fit85]. Then, \mathcal{T}_P is an approximation of T_P and the well-founded model of P satisfies $WF_P = WF(\mathcal{T}_P)$ [DMT00a]. Moreover, for weakly stratified programs, WF_P is two-valued [VRS91]. By Corollary 4.4

$$WF_P = WF(\mathcal{T}_P) \leq_p WF(U_P).$$

Since $WF(U_P)$ is consistent, and WF_P is complete, it follows that $WF_P = WF(U_P)$, as required. \square

We now show that in general, attractive properties of ultimate semantics come at a price. Namely, we have the following two theorems.

Theorem 5.3 *The problem “given a finite propositional logic program P , decide whether P has a complete ultimate stable model” is Σ_2^P -complete.*

Theorem 5.4 *The problems “given a finite propositional logic program, compute the ultimate well-founded fixpoint of P ” and “given a finite propositional logic program, compute the ultimate Kripke-Kleene fixpoint of P ” are in the class Δ_2^P .*

These results might put in doubt the usefulness of ultimate semantics. However, for wide classes of programs the complexity does not grow. Let k be a fixed integer. We define the class \mathcal{E}_k to consist of all logic programs P such that for every atom $p \in At(P)$ at least one of the following conditions holds:

1. P contains at most k clauses with p as the head;
2. the body of each clause with the head p consists of at most two elements;
3. the body of each clause with the head p contains at most one positive literal;
4. the body of each clause with the head p contains at most one negative literal.

Theorem 5.5 *The problem “given a finite propositional logic program from class \mathcal{E}_k , decide whether P has a complete ultimate stable model” is NP-complete.*

Theorem 5.6 *The problem “given a finite propositional logic program from class \mathcal{E}_k , compute the ultimate well-founded fixpoint of P ” is in P.*

We will now prove these results. If P is a finite propositional program, then it follows directly from the definition of the ultimate Kripke-Kleene fixpoint of T_P (that is, the ultimate Kripke-Kleene model of P) that it can be computed by means of polynomially many (in the size of P) evaluations of the operator $U_P(I, J)$, where $I \subseteq J$ are interpretations, with all other computational tasks taking only polynomial amount of time.

Let us also note that I is a complete ultimate stable model of P if and only if $I = lfp(U_P(\cdot, I))$. Thus, to verify whether I is a complete ultimate stable model, it is enough to iterate the operator $lfp(U_P(\cdot, I))$ starting with the empty interpretation. The number of iterations needed to reach the least fixpoint is again polynomial in the size of P with all other needed tasks taking polynomial time only. A similar discussion shows that the ultimate well-founded model of P can be computed by means of polynomially many evaluations of

the form $U_P(I, J)$.

It follows that evaluating $U_P(I, J)$, where $I \subseteq J$, is at the heart of computing the ultimate Kripke-Kleene, well-founded and complete stable models of a program P . Hence, we will now focus on this task.

Let P be a logic program and let p be an atom in P . For every rule $r \in P$ such that p is the head of r , we define B_r to be the conjunction of all literals in the body of r . For every atom p , we denote by $B_P(p)$ the disjunction of all formulas B_r , where r ranges over all rules in P with the head p . When p is the head of no rule in P then we set $B_P(p) = \perp$ (empty disjunction).

Every logic program P has a *normal* representation. It is the collection of rules $p \leftarrow B_P(p)$, where p ranges over all atoms of P . The definition of the operator T_P extends, in a straightforward way, to the case when P is given in its normal form defined above. Moreover, if P is a logic program and Q is its normal representation, $T_P = T_Q$. Thus, in the remainder of this section, without loss of generality we will assume that programs are given by means of their normal representations.

Let us recall that

$$U_P^1(I, J) = \text{glb}(T_P([I, J])) = \bigcap_{I \subseteq K \subseteq J} T_P(K)$$

and

$$U_P^2(I, J) = \text{lub}(T_P([I, J])) = \bigcup_{I \subseteq K \subseteq J} T_P(K).$$

Let I and J be two interpretations such that $I \subseteq J$. We define the *reduct* $P_{I,J}$ of P to be the program obtained from P by substituting in each body formula $B_P(p)$, any atom r by \mathbf{f} if $r \notin J$ and any atom r by \mathbf{t} if $r \in I$. Note that all body atoms of $P_{I,J}$ is an element of $J \setminus I$.

We have the following simple properties. An atom p of P belongs to $U_P^1(I, J)$ if and only if for every interpretation $K \in [\emptyset, J \setminus I]$, the formula $B_{P_{I,J}}(p)$ is true in K (or, equivalently, if and only if the formula $B_{P_{I,J}}(p)$ is a tautology). An atom p of P belongs to $U_P^2(I, J)$ if and only if for some interpretation $K \in [\emptyset, J \setminus I]$, the formula $B_{P_{I,J}}(p)$ is true in K (or, equivalently, if and only if the formula $B_{P_{I,J}}(p)$ is satisfiable).

From the second property it follows that computing $U_P^2(I, J)$ is easy — it can be accomplished in polynomial time (in the size of P). Indeed, since $B_{P_{I,J}}(p)$ is a DNF formula, its satisfiability can be decided in polynomial time and the claim follows. Thus, from now on we will focus on the task of computing $U_P^1(I, J)$.

The problem to decide whether a DNF formula is a tautology is co-NP-complete. Thus, the problem to compute the ultimate Kripke-Kleene and well-founded models of a program P is in the class Δ_2^P . Consequently, Theorem 5.4 follows.

It also follows that checking whether for an interpretation J , $J = \text{lfp}(U_P^1(\cdot, J))$ is in Δ_2^P . Hence, the problem to decide whether a program has a complete ultimate stable fixpoint is in the class Σ_2^P .

We will now show the Σ_2^P -hardness of the problem of existence of a complete ultimate stable model of a program P . Let φ be a propositional formula and let I be an interpretation (a set of atoms). We recall that the following problem is Σ_2^P -complete: Given a DNF formula φ over variables $x_1, \dots, x_m, y_1, \dots, y_n$, decide whether there is a truth assignment $I \subseteq \{x_1, \dots, x_m\}$ such that φ_I is a tautology, where φ_I is the formula obtained by replacing in φ all occurrences of atoms from I with \mathbf{t} , and by replacing all occurrences of atoms from $\{x_1, \dots, x_m\} \setminus I$ with \mathbf{f} .

We will reduce this problem to our problem. For each x_i , $i = 1, \dots, m$, in φ we introduce a new variable x'_i . We also introduce two new atoms p and q . By φ' we denote the formula obtained from φ by replacing literals $\neg x_i$ in the disjuncts of φ with new atoms x'_i . We define a program $P(\varphi)$ to consist of the following clauses:

1. $x_i \leftarrow \mathbf{not}(x'_i)$ and $x'_i \leftarrow \mathbf{not}(x_i)$, for every $i = 1, \dots, m$
2. $y_i \leftarrow \varphi'$, for every $i = 1, \dots, n$
3. $p \leftarrow \varphi'$
4. $q \leftarrow \mathbf{not}(p), \mathbf{not}(q)$.

We will show that there is $I \subseteq \{x_1, \dots, x_m\}$ such that φ_I is a tautology if and only if $P(\varphi)$ has an ultimate complete stable model.

It is easy to see that the following properties hold for every fixpoint M of $T_{P(\varphi)}$:

1. q is false in M (if q is true in M , $T_{P(\varphi)}$ does not derive q);
2. p is true in M (otherwise $T_{P(\varphi)}$ derives q);
3. y_1, \dots, y_n are true in M (since their rules have the same bodies as p);
4. for each x_i , either x_i or x'_i is true in M .

For a subset $I \subseteq \{x_1, \dots, x_m\}$, let us define $\bar{I} = I \cup \{x'_i : x_i \notin I\}$. It follows from the properties listed above that for each fixpoint M of $T_{P(\varphi)}$ and, a fortiori, if M is a complete ultimate stable model of $P(\varphi)$, there exists an I such that

$$M = \bar{I} \cup \{p, y_1, \dots, y_n\}.$$

Thus, it suffices to show that if $I \subseteq \{x_1, \dots, x_m\}$ then $M = \bar{I} \cup \{p, y_1, \dots, y_n\}$ is a complete ultimate stable model of $P(\varphi)$ if and only if φ_I is a tautology.

It is easy to verify that for every set $M = \bar{I} \cup \{p, y_1, \dots, y_n\}$ and for every $J \subseteq M$, $U^1(J, M)$ satisfies the following properties:

1. $U^1_{P(\varphi)}(J, M) \cap \{x_1, \dots, x_n, x'_1, \dots, x'_n\} = \bar{I}$
2. $U^1_{P(\varphi)}(J, M) \cap \{y_1, \dots, y_n, p, q\}$ is either \emptyset or $\{y_1, \dots, y_n, p\}$, since bodies of rules of y_1, \dots, y_n, p are identical.

Thus, we find that $U^1_{P(\varphi)}(J, M)$ is either \bar{I} or M and, consequently, $U^1_{P(\varphi)}(\cdot, M)$ has a least fixpoint, which is either \bar{I} or M . Hence $M = \bar{I} \cup \{p, y_1, \dots, y_n\}$ is a complete ultimate stable model of $P(\varphi)$ if and only if \bar{I} is not a fixpoint of $U^1_{P(\varphi)}(\cdot, M)$, that is if $U^1_{P(\varphi)}(\bar{I}, M) = M$. Consequently, all we need to prove is that $p \in U^1_{P(\varphi)}(\bar{I}, M)$ if and only if φ_I is a tautology.

Let us recall that $p \in U^1_{P(\varphi)}(\bar{I}, M)$ if and only if for every interpretation $K \in [\emptyset, M \setminus \bar{I}]$, the formula $B_{P(\varphi)_{\bar{I}, M}}(p)$ is true in K , that is, if and only if the formula $B_{P(\varphi)_{\bar{I}, M}}(p)$ is a tautology. Let us observe that $B_{P(\varphi)}(p) = \varphi'$. Thus, it is easy to see that $B_{P(\varphi)_{\bar{I}, M}}(p)$ is logically equivalent to φ_I . Consequently, the claim and Theorem 5.3 follows.

The problems of interest restricted to programs from the class \mathcal{E}_k become easier. Let us recall that the decision whether an atom $p \in At(P)$ belongs to $U^1_P(I, J)$ boils down to the decision whether the formula $B_{P_{I, J}}(p)$ is a tautology. If P is in the class \mathcal{E}_k , this question can be resolved in polynomial time. Thus, the ultimate Kripke-Kleene and the well-founded models for programs in \mathcal{E}_k can be computed in polynomial time. Thus, Theorem 5.6 follows.

Similarly, it takes only polynomial time to verify whether an interpretation I satisfies $I = lfp(U^1_P(\cdot, I))$. Thus, the problem to decide whether a program from \mathcal{E}_k has a complete ultimate stable model is in NP. To prove completeness, we observe that for purely negative programs:

1. there is no difference between complete stable fixpoints and complete ultimate stable fixpoints
2. purely negative programs are in \mathcal{E}_k
3. the problem of existence of complete stable fixpoints for purely negative programs is NP-complete.

Thus, Theorem 5.5 follows.

6 CONCLUSIONS AND DISCUSSION

We extended our algebraic framework [DMT00a, DMT00b] for studying semantics of nonmonotonic reasoning systems. The main contribution of this paper is the notion of an ultimate approximation. We argue that the Kripke-Kleene, well-founded and stable fixpoints of the ultimate approximation of an operator O can be regarded as the Kripke-Kleene, well-founded and stable fixpoints of the operator O itself. In earlier approaches, to study fixpoints of an operator O one needed to *select* an appropriate approximation operator. There were, however, no principled, algebraic ways to do so. In the present paper, we find a *distinguished* element in the space of all approximations and propose this particular approximation (ultimate approximation) to study fixpoints of O .

A striking feature of our approach is the ease with which it can be applied in any context where semantics emerge as fixpoints of operators. We applied this approach here in the context of logic programming and obtained a family of new semantics for logic programs: the ultimate Kripke-Kleene, the ultimate well-founded and the ultimate stable-model semantics. These semantics are well motivated and have attractive properties. First, they are preserved when we modify the program, as long as the 2-valued provability operator stays the same (the property that does not hold in general for standard semantics). Second, the ultimate Kripke-Kleene and the well-founded semantics are stronger (in general) than their standard counterparts, yet approximate the collection of all fixpoints of O and the collection of all stable fixpoints of O , respectively. The disadvantage is that their complexity is higher. But, as we noticed, for large classes of programs there is actually no loss in efficiency of computing ultimate semantics.

This approach can also be applied to default and autoepistemic logics and results in new semantics with appealing epistemological features². It was also recently used to define a precise semantics for logic programs with aggregates [DPB01].

We end this discussion with comments on a possible broader role of the approximation theory. One common concern when designing semantics of nonmonotonic logics is to avoid models justified by ungrounded or self-supporting (circular) arguments. The well-founded fixpoints (semantics) avoid such arguments. Groundedness is also a fundamental feature of induc-

²We will include a more extensive discussion of these applications in the journal version of the paper.

tion, a constructive way in which humans specify concepts both in commonsense reasoning settings and in formal considerations. In its simplest form induction relies only on positive information. In general, however, it may make references to negative information, too. In either form it is a nonmonotonic specification mechanism. As argued in [Den98], the well-founded semantics generalizes existing formalizations of induction (for instance, positive induction and iterated induction).

Acknowledgments

This material is based upon work supported by the National Science Foundation under Grants No. 9874764 and 0097278. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

References

- [AvE82] K.R. Apt and M.H. van Emden. Contributions to the theory of logic programming. *Journal of the ACM*, 29(3):841–862, 1982.
- [Cla78] K.L. Clark. Negation as failure. In H. Gallaire and J. Minker, editors, *Logic and data bases*, pages 293–322. Plenum Press, New York-London, 1978.
- [Den98] M. Denecker. The well-founded semantics is the principle of inductive definition. In J. Dix, L. Fariñas del Cerro, and U. Furbach, eds., *Logics in Artificial Intelligence*, LNAI, volume 1489, Springer-Verlag 1998.
- [DMT00a] M. Denecker, V. Marek, and M. Truszczyński. Approximations, stable operators, well-founded fixpoints and applications in nonmonotonic reasoning. In J. Minker, editor, *Logic-Based Artificial Intelligence*, pages 127–144. Kluwer Academic Publishers, 2000.
- [DMT00b] M. Denecker, V. Marek, and M. Truszczyński. Unified semantic treatment of default and autoepistemic logics. In *Principles of Knowledge Representation and Reasoning, Proceedings of the Seventh International Conference (KR2000)*, pages 74 – 84. Morgan Kaufmann Publishers, 2000.
- [DPB01] M. Denecker, N. Pelov, and M. Bruynooghe. Well-founded and stable semantics for logic programs with aggregates. In *Proceedings of ICLP-01*, LNCS 2237, Springer-Verlag, 2001.
- [Fit85] M. C. Fitting. A Kripke-Kleene semantics for logic programs. *Journal of Logic Programming*, 2(4):295–312, 1985.
- [Fit01] M. C. Fitting. Fixpoint semantics for logic programming – a survey. *Theoretical Computer Science*, 2002. To appear.
- [Gin88] M.L. Ginsberg. Multivalued logics: a uniform approach to reasoning in artificial intelligence. *Computational Intelligence*, 4:265–316, 1988.
- [GL88] M. Gelfond and V. Lifschitz. The stable semantics for logic programs. In R. Kowalski and K. Bowen, editors, *Proceedings of the 5th International Conference on Logic Programming*, pages 1070–1080. MIT Press, 1988.
- [Kun87] K. Kunen. Negation in logic programming. *Journal of Logic Programming*, 4(4):289–308, 1987.
- [Mar76] G. Markowsky. Chain-complete posets and directed sets with applications. *Algebra Universalis*, 6(1):53–68, 1976.
- [Moo84] R.C. Moore. Possible-world semantics for autoepistemic logic. In *Proceedings of the Workshop on Non-Monotonic Reasoning*, pages 344–354, 1984. Reprinted in: M. Ginsberg, ed., *Readings on non-monotonic reasoning*, pp. 137–142, Morgan Kaufmann, 1990.
- [Prz90] T.C. Przymusiński. The well-founded semantics coincides with the three-valued stable semantics. *Fundamenta Informaticae*, 13(4):445–464, 1990.
- [Rei80] R. Reiter. A logic for default reasoning. *Artificial Intelligence*, 13(1-2):81–132, 1980.
- [vEK76] M.H. van Emden and R.A. Kowalski. The semantics of predicate logic as a programming language. *Journal of the ACM*, 23(4):733–742, 1976.
- [VRS91] A. Van Gelder, K.A. Ross, and J.S. Schlipf. The well-founded semantics for general logic programs. *Journal of the ACM*, 38(3):620–650, 1991.